

A problem involving the p -Laplacian operator

Ratan K. Giri & D. Choudhuri*

Abstract

Using a variational technique we guarantee the existence of a solution to the *resonant Lane-Emden* problem $-\Delta_p u = \lambda|u|^{q-2}u$, $u|_{\partial\Omega} = 0$ if and only if a solution to $-\Delta_p u = \lambda|u|^{q-2}u + f$, $u|_{\partial\Omega} = 0$, $f \in L^{p'}(\Omega)$ (p' being the conjugate of p), exists for $q \in (1, p) \cup (p, p^*)$ under a certain condition for both the cases, i.e., $1 < q < p < p^*$ and $1 < p < q < p^*$ - the sub-linear and the super-linear cases.

keywords: p -laplacian; elliptic PDE; Palais-Smale condition; Sobolev space.

AMS classification: 35A15, 35A01.

1 Introduction

The study of partial differential equations involving a p -laplacian differential operator has become a major case of study in the recent times although it is still far from being completely understood, especially when $p = 1$ or ∞ . A few evidences of the limiting case can be found in [1], [2]. When $p = 2$, the usual Laplacian is obtained for which a vast literature exists ([4], [5] and the references therein). For $p \neq 2$ the p -Laplace operator has physical applications in the study of non-Newtonian fluids (dilatant fluids when $p > 2$) [6]. In practical life most of the problems are non linear by nature for which a numerical solution is sought for, however, unearthing the existence of solution leads to a rich theory hidden behind the partial differential equation. The problems we are going to address in this article are the following. Let Ω be a bounded subset of $\mathbb{R}^n$, $n \geq 3$ with a Lipschitz boundary $\partial\Omega$. Given $1 < p < \infty$ and $q \in (1, p) \cup (p, p^*)$, where $p^* = \frac{np}{n-p}$ if $1 < p < n$ and $p^* = \infty$ if $p \geq n$, we consider the following problems.

1. $-\Delta_p u = \lambda|u|^{q-2}u$, $u|_{\partial\Omega} = 0$. This problem is also known as the resonant Lane-Emden problem.
2. $-\Delta_p u = \lambda|u|^{q-2}u + f$, $f \in L^{p'}(\Omega)$, $u|_{\partial\Omega} = 0$.

where λ is a real number, $\Delta_p = \nabla \cdot (|\cdot|^{p-2} \nabla \cdot)$. Throughout this paper we shall refer the problems in 1 and 2 as the first and the second problem respectively.

*Corresponding author: dc.iit12@gmail.com

We call the first problem to be of sub-critical type if $1 < q < p < p^*$ and of super-critical type when $p^* > q > p > 1$. It is found in [7] that a unique solution exists to the first problem for the sub-critical case whereas uniqueness is lost for the super-critical case. Readers interested in knowing more about the first problem can refer to examples found in [8], [9], where the domain is ring shaped for $q \sim p^*$ and the solution is non-unique. Kawohl [10] showed the same but the domain which was considered is of annulus type with the annulus being sufficiently small in size. Dancer [11] showed that if $p = 2$ and Ω is a general domain then a unique solution exists to the first problem. Uniqueness is also guaranteed in [12] for the sub-linear case whereas a subdifferential method has been used to prove existence in [13] for both sub and super critical cases.

In this paper we will use a well known variational technique to show the existence of a solution in $W_0^{1,p}(\Omega) = \{v \in L^p(\Omega) : \nabla v \in L^p(\Omega), v|_{\partial\Omega} = 0\}$. A Fredholm type alternative is also proposed thus showing a connection between the first and the second problem. We organize the paper into two sections. In Section 2 we give the Mathematical formulation. In Section 3 we discuss a few preliminary results and the main result.

2 Mathematical formulation

The following definitions and theorems will be used in the main result we prove.

2.1 Definition: Let X be a Banach space and $H : X \rightarrow \mathbb{R}$ a C^1 functional. It is said to satisfy the *Palais-Smale condition* (PS) if the following holds:

Whenever $\{u_n\}$ is a sequence in X such that $\{H(u_n)\}$ is bounded and $H'(u_n) \rightarrow 0$ strongly in X' (the dual space), then $\{u_n\}$ has a strongly convergent subsequence.

The (PS) condition is a strong condition as very “well-behaved” function do not satisfy it (*Example:* $f(x) = c$, $x \in \mathbb{R}$, c a real constant).

We now state the following important theorem due to Ambrosetti and Rabinowitz [14] which is a common tool used in the theory of modern PDEs.

Mountain-pass theorem: Let $H : X \rightarrow \mathbb{R}$ be a C^1 functional satisfying (PS). Let $u_0, u_1 \in X$, $c_0 \in \mathbb{R}$ and $r > 0$ such that

1. $\|u_1 - u_0\| > r$
2. $H(u_0), H(u_1) < c_0 \leq H(v)$, $\forall v$ such that $\|v - u_0\| = r$. Then H has a critical value $c \geq c_0$ defined by

$$c = \inf_{\Gamma \in \wp} \max_{t \in [0,1]} H(\Gamma(t)) \quad (1)$$

where $\wp$ is the collection of all continuous paths $\Gamma : [0, 1] \rightarrow X$ such that $\Gamma(0) = u_0$, $\Gamma(1) = u_1$.

2.2 Weak formulation of the problem: We now give the weak formulation of the first problem. We say that $u \in W_0^{1,p}(\Omega)$ is a weak solution of the first problem if

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla v dx - \lambda \int_{\Omega} |u|^{q-2} u v dx = 0 \quad (2)$$

for every $v \in W_0^{1,p}(\Omega)$.

The weak solutions of the Lane-Emden problem are the critical points of the energy function defined by

$$J_q(u) = \frac{1}{p} \int_{\Omega} |\nabla u|^p dx - \frac{\lambda}{q} \int_{\Omega} |u|^q dx. \quad (3)$$

The following compact embedding theorems, due to Rellich-Kondrasov have been used in our work.

$$\begin{aligned} &\text{if } p < n, W_0^{1,p}(\Omega) \hookrightarrow L^q(\Omega), \quad 1 \leq q < p^*, \\ &\text{if } p = n, W_0^{1,p}(\Omega) \hookrightarrow L^q(\Omega), \quad 1 \leq q < \infty, \\ &\text{if } p > n, W_0^{1,p}(\Omega) \hookrightarrow C(\overline{\Omega}). \end{aligned}$$

We consider the non-homogeneous counterpart of the first problem - which is the second problem - and is as follows.

$$\begin{aligned} -\Delta_p u &= \lambda |u|^{q-2} u + f, \\ u|_{\partial\Omega} &= 0, \end{aligned} \quad (4)$$

where $f \in L^{p'}(\Omega)$, p' being the conjugate of p and is equal to $\frac{p}{p-1}$. Let the corresponding functional be denoted by J which is defined as follows.

$$J(u) = \frac{1}{p} \int_{\Omega} |\nabla u|^p dx - \frac{\lambda}{q} \int_{\Omega} |u|^q dx - \int_{\Omega} f u dx. \quad (5)$$

The Fréchet derivative of J , which is in $W_0^{-1,p'}(\Omega)$ where $p' = \frac{p}{p-1}$, is

$$\langle J'(u), v \rangle = \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla v dx - \lambda \int_{\Omega} |u|^{q-2} u v dx - \int_{\Omega} f v dx, \quad (6)$$

$\forall v \in W_0^{1,p}(\Omega)$. Thus $u \in W_0^{1,p}(\Omega)$ is a weak solution of the second problem if

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla v dx - \lambda \int_{\Omega} |u|^{q-2} u v dx - \int_{\Omega} f v dx = 0.$$

For the sake of further analysis we redefine the functional as follows.

$$J_q(u) = -\chi_{(1,p)}(q) J(u) + \chi_{(p,p^*)}(q) J(u), \quad (7)$$

where χ is the indicator function. From the sections which follow we shall use the functional in (7).

3 Few preliminary results and the main theorem

The main result of this paper is as follows. The problem $-\Delta_p u = \lambda|u|^{q-2}u$, $u|_{\partial\Omega} = 0$ has a weak solution if and only if the problem $-\Delta_p u = \lambda|u|^{q-2}u + f$, $u|_{\partial\Omega} = 0$, where $f \in L^{p/p-1}(\Omega)$, has a weak solution. We prove the result for $p < n$. The case of $p \geq n$ follows the same proof as in the case $p < n$ which is based on the results on compact embedding stated after equation (3). But first we present a few technical lemmas on which the proof of this result relies upon.

We first assume that a solution exists to the problem

$$\begin{aligned} -\Delta_p u &= \lambda|u|^{q-2}u, \\ u|_{\partial\Omega} &= 0. \end{aligned} \tag{8}$$

Theorem 3.1. *The mapping J_q defined in (7) is a C^1 -functional over $W_0^{1,p}(\Omega)$.*

Proof. We first prove that the functional J' is continuous which will imply that J'_q is continuous and hence the theorem will follow. Consider

$$\begin{aligned} | \langle J'(u), v \rangle | &\leq \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla v dx + |\lambda| \int_{\Omega} |u|^{q-1} v dx + \int_{\Omega} |f| |v| dx \\ &\leq \|\nabla u\|_{\frac{p}{p-1}} \|\nabla v\|_p + |\lambda| \|u\|_{\frac{q}{q-1}} \|v\|_q + \|f\|_{\frac{p}{p-1}} \|v\|_p \\ &\leq \left[\|\nabla u\|_{\frac{p}{p-1}} + C_1 |\lambda| \|u\|_{\frac{q}{q-1}} + C_2 \|f\|_{\frac{p}{p-1}} \right] \|\nabla v\|_p, \quad \forall v \in W_0^{1,p}(\Omega), \end{aligned} \tag{9}$$

where C_1, C_2 are the constants due to the embedding of $W_0^{1,p}(\Omega)$ in $L^q(\Omega)$ for $q \in [1, p^*]$. From (8)&(9) one can see that J is a C^1 functional over $W_0^{1,p}(\Omega)$. $\square$

Theorem 3.2. *There exists $u_0, u_1 \in W_0^{1,p}(\Omega)$ and a positive real number c_0 such that $J_q(u_0), J_q(u_1) < c_0$ and $J_q(v) \geq c_0$, for every v satisfying $\|v - u_0\|_{1,p} = r$.*

Proof. Let $u_0 = 0$. Clearly u_0 is a solution of (8) and $J_q(0) = 0$. Now let $w \in B(0, 1)$ in $W_0^{1,p}(\Omega)$ and consider $v = u_0 + rw$ for $r > 0$ and hence $\|v - u_0\|_{1,p} = r$. We first show the existence of r such that $\|v - u_0\|_{1,p} = r_0$ and for which $J(v) \geq c_0$ for each $v \in B(0, r_0)$.

Let $p < q < p^*$. Now

$$\begin{aligned} J_q(u_0 + rw) - J_q(u_0) &= \frac{r^p}{p} \int_{\Omega} |\nabla w|^p dx - \frac{r^q \lambda}{q} \int_{\Omega} |w|^q dx - r \int_{\Omega} f w dx, \\ &= \frac{r^p}{p} - \frac{r^q \lambda}{q} \int_{\Omega} |w|^q dx - r \int_{\Omega} f w dx. \end{aligned} \tag{10}$$

Further, $|w|_{1,p} = 1$ and hence $|\int_{\Omega} w^p dx| \leq \int_{\Omega} |w|^p dx \leq c \|w\|_p^p \leq c_1 |w|_{1,p} = c_1$. Similarly, $|\int_{\Omega} w^q dx| \leq c_2$. Using these arguments leads to

$$\begin{aligned} J_q(u_0 + rw) - J_q(u_0) &\geq r \left[\frac{r^{p-1}}{p} - \frac{r^{q-1} \lambda}{q} c_2 - c_1^{1/p} \|f\|_{p'} \right], \\ &= c'. \end{aligned} \tag{11}$$

We first analyze the term $\left[\frac{r^{p-1}}{p} - \frac{r^{q-1}\lambda}{q}c_2 - c_1^{1/p}\|f\|_{p'} \right] = F(r)$ (say). Clearly $F(0) < 0$ and for $r_0 = \left(\frac{q(p-1)}{p(q-1)} \frac{1}{\lambda c_2} \right)^{\frac{1}{q-p}}$ we see that $F'(r_0) = 0$. A bit of calculus guarantees that $F''(r_0) < 0$ and hence r_0 is a maximizer of F . If $0 < \lambda < \lambda_1 = \frac{q(p-1)}{p(q-1)} \cdot \left(\frac{p(q-1)}{q-p} \cdot c_1^{\frac{1}{p}} \|f\|_{p'} \right)^{\frac{1}{1-p}}$ then $F(r_0) > 0$. As $r \rightarrow \infty$ we have $F(r) \rightarrow -\infty$. Hence there exists $r_1, r_2 > 0$ and $r_1 < r_0 < r_2$ such that $F(r) > 0$ for each $r \in (r_1, r_2)$. We choose $r = r_0$ such that $\|v - u_0\|_{1,p} = r_0$ and for which $J_q(v) \geq c'$ for each $v \in B(0, r_0)$. Similarly, if $1 < q < p$ then according to the definition of J_q we now have

$$\begin{aligned} J_q(u_0 + rw) - J_q(u_0) &= -J(u_0 + rw) + J(u_0) \\ &\geq r \left[-\frac{r^{p-1}}{p} + \frac{r^{q-1}\lambda}{q}c_2 + c_1^{1/p}\|f\|_{p'} \right], \\ &= c''. \end{aligned} \tag{12}$$

Using the same argument as for the case of $p < q < p^*$ we find r and $0 < \lambda < \lambda_2$ such that $J_q(v) \geq c''$ for all $\|v - u_0\| = r$. We choose $\lambda' = \min\{\lambda_1, \lambda_2\}$ such that $0 < \lambda < \lambda'$ and $c_0 = \min\{c', c''\}$.

Choice of u_1 : Let w_p be the first eigen vector of $-\Delta_p$, i.e., $-\Delta_p w_p = \lambda_p |w_p|^{p-1} w_p$, where λ_p is the first eigen value of $-\Delta_p$. The first eigen value of the p -laplacian operator is strictly positive [3]. Consider the function $g = kw_p$, $k \in \mathbb{R}$, $\|w_p\|_{1,p} = 1$ and $p < q < p^*$. Note that,

$$J_q(g) = \left(\frac{k^p}{p} - \frac{\lambda k^q \int_{\Omega} |w_p|^q dx}{q} \right) - kC,$$

where $C = \int_{\Omega} f w_p dx$. Since $p < q < p^*$, we observe k can be chosen arbitrarily large so that $\frac{k^p}{p} - \frac{\lambda k^q \int_{\Omega} |w_p|^q dx}{q} - k_0 C < 0$. Then $J_q(kw_p) < 0$ and hence $J_q(kw_p) < J_q(u_0)$. Thus we can choose $u_1 = k_0 w_p$, where $k_0 > r_0$. Then $\|u_1 - u_0\|_{1,p} > r_0$. Similarly for $1 < q < p$ we have

$$J_q(g) = \left(-\frac{k^p}{p} + \frac{\lambda k^q \int_{\Omega} |w_p|^q dx}{q} \right) + kC,$$

and k can be chosen large enough to make $J_q(g) < 0$. Hence the result. $\square$

Theorem 3.3. J_q satisfies the Palais-Smale condition.

Proof. Let us consider the case for which $p < q < p^*$. The other case for $1 < q < p$ follows similarly. Let u_n be a sequence in $W_0^{1,p}(\Omega)$ such that $|J_q(u_n)| \leq M$ and $J'_q(u_n) \rightarrow 0$ as $n \rightarrow \infty$ in $W_0^{-1,p'}(\Omega)$, p' being the conjugate of p . Now

$$J_q(u_n) = \frac{1}{p} \int_{\Omega} |\nabla u_n|^p dx - \frac{\lambda}{q} \int_{\Omega} |u_n|^q dx - \int_{\Omega} f u_n dx, \tag{13}$$

$$\begin{aligned} \langle J'_q(u_n), v \rangle &= \int_{\Omega} |\nabla u_n|^{p-2} \nabla u_n \cdot \nabla v dx - \lambda \int_{\Omega} |u_n|^{q-2} u_n v dx - \int_{\Omega} f v dx, \forall v \in W_0^{1,p}(\Omega). \end{aligned} \tag{14}$$

Consider the following.

$$\langle J'_q(u_m), u_m \rangle = \int_{\Omega} |\nabla u_m|^p dx - \lambda \int_{\Omega} |u_m|^q dx - \int_{\Omega} f u_m dx, \quad (15)$$

$$\begin{aligned} J_q(u_m) &= \frac{1}{p} \int_{\Omega} |\nabla u_m|^p dx - \frac{\lambda}{q} \int_{\Omega} |u_m|^q dx - \int_{\Omega} f u_m dx, \\ &= \frac{1}{p} |u_m|_{1,p}^p - \frac{\lambda}{q} \int_{\Omega} |u_m|^q dx - \int_{\Omega} f u_m dx \\ \lambda \int_{\Omega} |u_m|^q dx &= \frac{q}{p} |u_m|_{1,p}^p - q J_q(u_m) - q \int_{\Omega} f u_m dx, \\ \frac{p-q}{p} |u_m|_{1,p}^p &= \langle J'_q(u_m), u_m \rangle - q J_q(u_m) - q \int_{\Omega} f u_m dx. \end{aligned} \quad (16)$$

This implies that $|u_m|_{1,p}$ is bounded. The above inequality in (14) clearly shows that u_n is bounded in $W_0^{1,p}(\Omega)$ and hence by Eberlein-Šmulian's theorem (refer Dunford-Schwartz [1; p. 430] [15]) it has a *weakly* convergent subsequence, say u_{n_k} , in $W_0^{1,p}(\Omega)$.

Claim. The sequence $\{u_{n_k}\}$ is strongly convergent in $W_0^{1,p}(\Omega)$.

Proof. Applying limit $k \rightarrow \infty$ to (14) (refer Appendix) and using the strong convergence of (u_{n_k}) in $L^q(\Omega)$ due to compact embedding we obtain

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla v dx = \lambda \int_{\Omega} |u|^{q-2} u v dx + \int_{\Omega} f v dx, \quad (17)$$

and we pass on the limit to (15) we get

$$\lim_{n \rightarrow \infty} |u_{n_k}|_{1,p}^p = \lambda \int_{\Omega} |u|^q dx + \int_{\Omega} f u dx = |u|_{1,p}^p. \quad (18)$$

Thus a weakly convergent sequence which is convergent in norm is strongly convergent. Hence $u_{n_k} \rightarrow u$ in $W_0^{1,p}(\Omega)$ as $k \rightarrow \infty$. $\square$

Thus by the Mountain-pass theorem an extreme point for H exists in $W_0^{1,p}(\Omega)$

We summarize the results proved in Theorems 3.1, 3.2 and 3.3 in the form of a theorem as follows.

Theorem 3.4. *Suppose $-\Delta_p u = \lambda |u|^{q-2} u$, $u|_{\partial\Omega} = 0$ has a solution. Then*

1. *the functional $J_q = -\chi_{(1,p)}(q)J(u) + \chi_{(p,p^*)}(q)J(u)$ where $J(u) = \frac{1}{p} \int_{\Omega} |\nabla u|^p dx - \frac{\lambda}{q} \int_{\Omega} |u|^q dx - \int_{\Omega} f u dx$ is C^1 and satisfies the Palais-Smale condition,*
2. *J_q satisfies the hypothesis of the Mountain-Pass theorem.*

Therefore J_q has an extreme point in $W_0^{1,p}(\Omega)$. In other words $-\Delta_p u = \lambda |u|^{q-2} u + f$, $f \in L^{p'}(\Omega)$, $u|_{\partial\Omega} = 0$ has a solution whenever $\lambda \in (0, \lambda']$ where $\lambda' = \min\{\lambda_1, \lambda_2\}$ as found in Theorem 3.2.

Conversely, suppose a solution to the problem

$$\begin{aligned} -\Delta_p u &= \lambda |u|^{q-2} u + f, \quad f \in L^{p'}(\Omega), \\ u|_{\partial\Omega} &= 0. \end{aligned} \quad (19)$$

We subdivide this situation into two different cases - namely, $1 < q < p$ (the sub-linear case) and $1 < p < q < p^*$ (the super-linear cases). Let $(f_n) \subset L^{p'}(\Omega)$ be a sequence such that $f_n \rightarrow 0$ in $L^{p'}(\Omega)$. By the assumption, to each f_n there exists a solution, say u_n .

We have $q \in (1, p) \cup (p, p^*)$ and

$$\begin{aligned} B[u, v] &= \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla v dx - \lambda \int_{\Omega} |u|^{q-2} u v dx, \\ &= \int_{\Omega} f v dx, \quad \forall v \in W_0^{1,p}(\Omega) \end{aligned} \quad (20)$$

where B is a ‘non linear form’ in two variables u and v . It is easy to check that $B(.,.)$ is the Fréchet derivative of the C^1 functional $\frac{1}{p} \int_{\Omega} |\nabla u|^p - \frac{\lambda}{q} \int_{\Omega} |u|^q$ and hence is continuous.

Clearly, for each $v \in W_0^{1,p}(\Omega)$ we have

$$\begin{aligned} B[u_n, v] &= \int_{\Omega} |\nabla u_n|^{p-2} \nabla u_n \cdot \nabla v dx - \lambda \int_{\Omega} |u_n|^{q-2} u_n v dx, \\ &= \int_{\Omega} f_n v dx, \\ &\leq \|f_n\|_{p'} \|v\|_p \rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned} \quad (21)$$

Hence $\int_{\Omega} f_n v dx \rightarrow 0$ as $n \rightarrow \infty$. Consider $T_n(v) = \int_{\Omega} |\nabla u_n|^{p-2} \nabla u_n \cdot \nabla v dx$. Then T_n is bounded linear over $W_0^{1,p}(\Omega)$ and $\|T_n\| = \| |\nabla u_n|^{p-1} \|_{p'}$. From the above definition of T_n , for a fixed $v \in W_0^{1,p}(\Omega)$ we have the sequence $(T_n(v))$ to be bounded which implies that $(T_n(v))$ is pointwise bounded. Thus by the uniform boundedness principle $(\|T_n\|)$ is bounded. Thus $\| |\nabla u_n|^{p-1} \|_{p'}$ is bounded. Hence, there exists a subsequence (u_{n_k}) which weakly converges to u_{∞} with respect to the $\|\cdot\|_{1,p}$ in $W_0^{1,p}(\Omega)$. Hence we have

$$\begin{aligned} \lim_{k \rightarrow \infty} \int_{\Omega} |\nabla v|^{p-2} \nabla v \cdot \nabla u_{n_k} dx &= \int_{\Omega} |\nabla v|^{p-2} \nabla v \cdot \nabla u_{\infty} dx, \quad \forall v \in W_0^{1,p}(\Omega). \\ \Rightarrow \lim_{k \rightarrow \infty} \int_{\Omega} |\nabla u_{n_l}|^{p-2} \nabla u_{n_l} \cdot \nabla u_{n_k} dx &= \int_{\Omega} |\nabla u_{n_l}|^{p-2} \nabla u_{n_l} \cdot \nabla u_{\infty} dx, \end{aligned} \quad (22)$$

for a fixed l . Therefore, since $u_{n_k} \rightharpoonup u_{\infty}$ in $W_0^{1,p}(\Omega)$ implies that $|\nabla u_{n_k}|^{p-1} \rightharpoonup |\nabla u_{\infty}|^{p-1}$ (for a subsequence) in $L^{p'}(\Omega)$ (Refer Appendix). But $W_0^{1,p}(\Omega) \hookrightarrow L^p(\Omega) \hookrightarrow W^{-1,p'}(\Omega)$ and hence

$$\begin{aligned} \lim_{l \rightarrow \infty} \int_{\Omega} |\nabla u_{n_l}|^{p-2} \nabla u_{n_l} \cdot \nabla v dx &= \int_{\Omega} |\nabla u_{\infty}|^{p-2} \nabla u_{\infty} \cdot \nabla v dx, \quad \forall v \in W_0^{1,p}(\Omega), \\ \Rightarrow \lim_{l \rightarrow \infty} \int_{\Omega} |\nabla u_{n_l}|^{p-2} \nabla u_{n_l} \cdot \nabla u_{\infty} dx &= \int_{\Omega} |\nabla u_{\infty}|^p dx. \end{aligned} \quad (23)$$

Hence, $\lim_{k \rightarrow \infty} \int_{\Omega} |\nabla u_{n_k}|^p dx = \int_{\Omega} |\nabla u_{\infty}|^p dx$. It immediately can be concluded that there exists a u_{∞} such that $u_n \rightarrow u_{\infty}$ in $W_0^{1,p}(\Omega)$. Hence using the continuity of $B[.,.]$ in (20) we have

$$\begin{aligned} \lim_{n \rightarrow \infty} B[u_n, v] &= \lim_{n \rightarrow \infty} \int_{\Omega} |\nabla u_n|^{p-2} \nabla u_n \cdot \nabla v dx - \lim_{n \rightarrow \infty} \lambda \int_{\Omega} |u_n|^{q-2} u_n v dx, \\ &= \lim_{n \rightarrow \infty} \int_{\Omega} f_n v dx, \\ \Rightarrow B[u_{\infty}, v] &= 0, \forall v \in W_0^{1,p}(\Omega). \end{aligned}$$

In other words

$$\int_{\Omega} |\nabla u_{\infty}|^{p-2} \nabla u_{\infty} \cdot \nabla v dx - \lambda \int_{\Omega} |u_{\infty}|^{q-2} u_{\infty} v dx = 0, \forall v \in W_0^{1,p}(\Omega). \quad (24)$$

We summarize the result proved as follows.

Theorem 3.5. *Suppose $-\Delta_p u = \lambda|u|^{q-2}u + f$, $f \in L^{p'}(\Omega)$, $u|_{\partial\Omega} = 0$ has a solution. If u_n is a solution of the PDE corresponds to f_n , where $f_n \in L^{p'}(\Omega)$ such that $f_n \rightarrow 0$ in $L^{p'}(\Omega)$, then we have $B[u_{\infty}, v] = 0$ for each $v \in W_0^{1,p}(\Omega)$ and thus u_{∞} is a solution to $-\Delta_p u = \lambda|u|^{q-2}u$, $u|_{\partial\Omega} = 0$.*

4 Appendix

We show that

$$\lim_{n \rightarrow \infty} \int_{\Omega} |\nabla u_n|^{p-2} \nabla u_n \cdot \nabla v dx = \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla v dx, \quad \forall v \in W_0^{1,p}(\Omega). \quad (25)$$

We divide the explanation into two cases:

Case 1: When $p > 2$.

This implies that p' , the conjugate of p , should be lesser than 2, i.e., $1 < p' < 2 < p$. Thus we have $W_0^{1,p}(\Omega) \hookrightarrow_{compact} L^{p'}(\Omega)$ (since $W_0^{1,p}(\Omega) \hookrightarrow_{compact} L^q(\Omega)$ for $q \in [1, p^*)$).

Since ∇u_n converges weakly to, say ∇u , in $L^p(\Omega)$, hence $\langle |\nabla u_n| - |\nabla u|, v \rangle \rightarrow 0$ for each $v \in L^{p'}(\Omega)$. Thus $\langle |\nabla u_n| - |\nabla u|, |\nabla u_n| - |\nabla u| \rangle \rightarrow 0$, i.e., $\|\nabla u_n\|_2 \rightarrow \|\nabla u\|_2$. Hence $\|\nabla u_n\|_{p'} \rightarrow \|\nabla u\|_{p'}$ because $p' < 2 < p$. By the Riesz-Fischer theorem [16], there exists a subsequence of ∇u_n which converges pointwise a.e., i.e., $|\nabla u_n(x)| \rightarrow |\nabla u(x)|$. So $|\nabla u_n(x)|^{p-1} \rightarrow |\nabla u(x)|^{p-1}$ and hence $|\nabla u_n|^{p-1} \rightharpoonup |\nabla u|^{p-1}$ in $L^{p'}(\Omega)$. Thus we have $\lim_{n \rightarrow \infty} \int_{\Omega} |\nabla u_n|^{p-2} \nabla u_n \cdot \nabla v dx = \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla v dx, \forall v \in W_0^{1,p}(\Omega)$.

Case 2: When $p < 2$.

This implies that p' , the conjugate of p , should be greater than 2, i.e., $p < 2 < p'$.

Look at the map $F : W_0^{1,p}(\Omega) \rightarrow L^{p'}(\Omega)$ defined by $u \mapsto |\nabla u|^{p-1}$. Consider the range of F , i.e., $R(F) = \{|\nabla u|^{p-1} : u \in W_0^{1,p}(\Omega)\}$.

Observe that the map F is bounded in the sense that bounded sets are mapped to bounded

sets. Hence if $u_n \rightharpoonup u$ in $W_0^{1,p}(\Omega)$ implies that (u_n) is bounded in $W_0^{1,p}(\Omega)$. Hence $(F(u_n)) = (|\nabla u_n|^{p-1})$ is bounded in $L^{p'}(\Omega)$. Since $L^{p'}(\Omega)$ is reflexive, hence there exists a subsequence of $|\nabla u_n|^{p-1}$ which weakly converges to, say, w in $L^{p'}(\Omega)$.

We have the following.

$u_n \rightharpoonup u$ in $W_0^{1,p}(\Omega)$ so $|\nabla u_n|^{p-1} \rightharpoonup w$ in $L^{p'}(\Omega)$. This implies that

$$\langle |\nabla u_n|^{p-1} - w, v \rangle \rightarrow 0, \forall v \in L^p(\Omega)$$

Since $p < 2 < p'$ hence $|\nabla u_n|^{p-1} - w \in L^p(\Omega)$. Thus $\| |\nabla u_n|^{p-1} - w \|_2 \rightarrow 0$ and hence $\| |\nabla u_n|^{p-1} - w \|_p \rightarrow 0$. Therefore we have a subsequence of $(|\nabla u_n|^{p-1})$ such that $|\nabla u_n|^{p-1} \rightarrow w$ pointwise a.e. (implying $|\nabla u_n| \rightarrow w^{\frac{1}{p-1}}$ pointwise a.e.) and so $|\nabla u_n| \rightharpoonup w^{\frac{1}{p-1}}$ in $L^p(\Omega)$. Hence $w = |\nabla u|^{p-1}$.

Thus in all the above cases we found the following.

$$\lim_{n \rightarrow \infty} \int_{\Omega} |\nabla u_n|^{p-2} \nabla u_n \cdot \nabla v dx = \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla v dx, \quad \forall v \in W_0^{1,p}(\Omega). \quad (26)$$

Hence by the compact embedding due to Rellich-Kondrachov it can be concluded $u_n \rightarrow u$ in $L^q(\Omega)$. Thus we also have

$$\lim_{n \rightarrow \infty} \int_{\Omega} |u_n|^{q-2} u_n \cdot \nabla v dx = \int_{\Omega} |u|^{q-2} u \cdot \nabla v dx, \quad \forall v \in W_0^{1,p}(\Omega). \quad (27)$$

5 Conclusions

The resonant Lane-Emden problem has been studied. An existence result has been established to the non-homogeneous Lane-Emden problem for the sub-linear - $1 < q < p < p^*$ and the super-linear case - $1 < p < q < p^*$ for $\lambda \in (0, \lambda']$ - λ' being sufficiently large - if it is assumed that a non-trivial solution exists to the homogeneous Lane-Emden problem for the sub-linear - $1 < q < p < p^*$ and the super-linear case - $1 < p < q < p^*$, which is basically an eigen value problem. We further proved the ‘converse’ that if the non-homogeneous problem has a solution then a solution to the homogeneous problem exists for both the sub and the super critical cases.

6 Acknowledgement

One of the authors (RKG) thanks the financial assistantship received from the Ministry of Human Resource Development (M.H.R.D.).

References

- [1] T. BHATTACHARYA, E.D. BENEDETTO and J. MANFREDI, Limits as $p \rightarrow \infty$ of $\Delta_p u = f$ and related extremal problems, *Rend. Sem. Mat. Univ. Politec. Torino*, special issue (1991), 15-68.

- [2] P. JUNTINEN, P. LINDQVIST, On the higher eigen values for the ∞ -eigen value problem, *Calc. Var. partial differential equations* **23** (2005), 169-192.
- [3] IDRISSE LY, The first eigen value for the p-laplacian operator, *Journal of inequalities of pure and applied Math.*, **6** (3) (2005), 1-28.
- [4] L.C. EVANS, *Partial Differential Equations* (Amer. Math. Soc., 2009).
- [5] KESAVAN, *Functional Analysis and applications* (New age international pvt. ltd., 2003).
- [6] NIKOS E. MASTORAKIS, HASSAN FATHABADI, On the solution of p-Laplacian for non-newtonian fluid flow, *W.S.E.A.S. transactions on Math.*, **8** (6) (2009), 238-245.
- [7] GREY ERCOLE, On the resonant Lane-Emden problem for the p-Laplacian, *Communications in Contemporary Mathematics*, **16** (4) (2014), 1350033-1-22.
- [8] P. DRÁBEK, A note on the nonuniqueness for some quasilinear eigenvalue problem, *Appl. Math. Lett.*, **13** (2000), 39-41.
- [9] J. GARCÍA AZORERO and I. PERAL ALONSO, On limits of solutions of elliptic problems with nearly critical exponent, *Comm. Partial Differential Equations*, **17**(1992), 2113-2126.
- [10] B. KAWOHL, Symmetry results for functions yielding best constants in Sobolev-type inequalities, *Discrete Contin. Dynam. Systems*, **6**(2000), 683-690.
- [11] E.N. DANCER, Real analyticity and non-degeneracy, *Math. Ann.*, **325**(2003), 369-392.
- [12] J.I. DÍAZ and J.E. SAA, Existence et unicité de solutions positives pour certaines équations elliptiques quasilineaires, *C.R. Acad. Sci. Paris*, **305**(1987), 521-524.
- [13] M. ÔTANI, Existence and nonexistence of nontrivial solutions of some nonlinear degenerate elliptic equations, *J. Funct. Anal.*, **76**(1988), 140-159.
- [14] A. AMBROSETTI and P.H. RABINOWITZ, Dual Variational Methods in Critical Point Theory and Applications, *J. Funct. Anal.*, **14**(1973), 349-381.
- [15] N. DUNFORD and J. SCHWARTZ, *Linear Operators. Part 2: Spectral theory*, Wiley (Interscience), New York, 1958.
- [16] G. BACHMAN and L. NARICI, *Functional Analysis*, Dover Publications, Mineola, New York, 1966.

RATAN KUMAR GIRI and D. CHOUDHURI

Department of Mathematics,

National Institute of Technology Rourkela, Rourkela - 769008, India